

A particular matrix and its relationship with Fibonacci numbers

Mario Catalani
 Department of Economics
 Via Po 53, 10124 Torino, Italy
 e-mail mario.catalani@unito.it

Abstract

Determinants and symmetric functions of the eigenvalues of matrices characterizing stochastic processes with independent increments. Relationships with Fibonacci numbers are derived.

1 Introduction

Let us consider the $n \times n$ symmetric matrix $\mathbf{A}_n$

$$\mathbf{A}_n = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & 2 & 2 & \cdots & 2 \\ 1 & 2 & 3 & \cdots & 3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 2 & 3 & \cdots & n \end{bmatrix},$$

that is

$$a_{ij} = \min(i, j), \quad i, j = 1, 2, \dots, n$$

This matrix is, up to a positive scalar, the covariance matrix of a stochastic process with increments which possess the same variance and are uncorrelated.

For future reference consider also the $(n - k + 1) \times (n - k + 1)$ matrix $\mathbf{C}_{n,k}$

$$\mathbf{C}_{n,k} = \begin{bmatrix} k & k & k & \cdots & k \\ k & k+1 & k+1 & \cdots & k+1 \\ k & k+1 & k+2 & \cdots & k+2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ k & k+1 & k+2 & \cdots & n \end{bmatrix}, \quad k \text{ integer, } 1 < k < n.$$

For general reference on matrix theory see, for example, [1], [2], [4].

2 Determinants

Let us consider the following determinants (dimensions $n \times n$)

$$\Delta_n(i_1, i_2, \dots, i_n) = \begin{vmatrix} i_1 & i_1 & i_1 & \cdots & i_1 \\ i_1 & i_1 + i_2 & i_1 + i_2 & \cdots & i_1 + i_2 \\ i_1 & i_1 + i_2 & i_1 + i_2 + i_3 & \cdots & i_1 + i_2 + i_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ i_1 & i_1 + i_2 & i_1 + i_2 + i_3 & \cdots & i_1 + i_2 + \cdots + i_n \end{vmatrix},$$

and

$$\begin{aligned} \Theta_n(i_1, i_2, \dots, i_n, i_{n+1}) &= \\ &= \begin{vmatrix} i_1 & i_1 + i_2 & i_1 + i_2 & \cdots & i_1 + i_2 \\ i_1 & i_1 + i_2 + i_3 & i_1 + i_2 + i_3 & \cdots & i_1 + i_2 + i_3 \\ i_1 & i_1 + i_2 + i_3 & i_1 + i_2 + i_3 + i_4 & \cdots & i_1 + i_2 + i_3 + i_4 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ i_1 & i_1 + i_2 + i_3 & i_1 + i_2 + i_3 + i_4 & \cdots & i_1 + i_2 + \cdots + i_{n+1} \end{vmatrix}, \end{aligned}$$

where $i_j, \forall j$ are integers ≥ 1 . Let us note that the determinant of $\mathbf{A}_n$ is $\Delta_n(1, 1, \dots, 1)$ and that of the matrix $\mathbf{C}_{n,k}$ is $\Delta_n(k, 1, \dots, 1)$.

We are going to prove that

$$\Delta_n(i_1, i_2, \dots, i_n) = \prod_{j=1}^n i_j, \quad (1)$$

and

$$\Theta_n(i_1, i_2, \dots, i_n, i_{n+1}) = i_1 \prod_{j=3}^{n+1} i_j. \quad (2)$$

We are going to prove the claim using induction.

First of all

$$\begin{aligned} \Delta_2(i_1, i_2) &= \begin{vmatrix} i_1 & i_1 \\ i_1 & i_1 + i_2 \end{vmatrix} \\ &= i_1(i_1 + i_2 - i_1) \\ &= i_1 i_2, \end{aligned}$$

and

$$\begin{aligned}
\Theta_2(i_1, i_2, i_3) &= \begin{vmatrix} i_1 & i_1 + i_2 \\ i_1 & i_1 + i_2 + i_3 \end{vmatrix} \\
&= i_1(i_1 + i_2 + i_3 - i_1 - i_2) \\
&= i_1 i_3.
\end{aligned}$$

Now assume that the claim holds for $n-1$. If we evaluate $\Delta_n(i_1, i_2, \dots, i_n)$ expanding with respect to the elements of the first row we have that all the algebraic complements except those relative to the first two elements of the row are equal to zero: indeed all the above have identical the first two columns, and we obtain

$$\begin{aligned}
\Delta_n(i_1, i_2, \dots, i_n) &= i_1 \Delta_{n-1}(i_1 + i_2, i_3, \dots, i_n) - i_1 \Theta_{n-1}(i_1, i_2, \dots, i_n) \\
&= i_1(i_1 + i_2)i_3 \cdots i_n - i_1 i_1 i_3 \cdots i_n \\
&= i_1 i_3 i_4 \cdots i_n (i_1 + i_2 - i_1) \\
&= \prod_{j=1}^n i_j.
\end{aligned}$$

Now reverting to $\Theta_n(i_1, i_2, \dots, i_n, i_{n+1})$ the same reasoning as before leads to

$$\begin{aligned}
\Theta_n(i_1, i_2, \dots, i_n, i_{n+1}) &= i_1 \Delta_{n-1}(i_1 + i_2 + i_3, i_4, \dots, i_n, i_{n+1}) \\
&\quad - (i_1 + i_2) \Theta_{n-1}(i_1, i_2 + i_3, i_4, \dots, i_{n+1}) \\
&= i_1(i_1 + i_2 + i_3)i_4 \cdots i_{n+1} - (i_1 + i_2)i_1 i_4 \cdots i_{n+1} \\
&= i_1 i_4 \cdots i_{n+1} (i_1 + i_2 + i_3 - i_1 - i_2) \\
&= i_1 \prod_{j=3}^{n+1} i_j.
\end{aligned}$$

This completes the proof.

As a corollary we see that immediately we have $|\mathbf{A}_n| = 1$, and $|\mathbf{C}_{n,k}| = k$.

3 Symmetric Functions of Eigenvalues

Just to fix notation, given n numbers

$$\{a_i\}_1^m = \{a_1, a_2, \dots, a_n\}$$

let $S_k^n \langle a_i \rangle$ denote the symmetric functions, that is

$$S_k^n \langle a_i \rangle = \sum_{\substack{i_1, \dots, i_k \\ i_1 < i_2 < \dots < i_k}} a_{i_1} a_{i_2} \cdots a_{i_k}, \quad 1 \leq k \leq n.$$

Now given a $n \times n$ matrix $\mathbf{A}$ with eigenvalues $\{\lambda_i\}$ we have

$$S_k^n \langle \lambda_i \rangle = \sum_i \left| \mathbf{A}_{(i)}^{(n-k)} \right|, \quad (3)$$

where

$$\mathbf{A}_{(i)}^{(h)}, \quad h = 0, 1, \dots, n-1$$

denotes one of the principal submatrices of $\mathbf{A}$ obtained deleting h rows and the corresponding h columns. The number of these submatrices is

$$\binom{n}{h}.$$

Note that the submatrices in Equation 3 have dimensions $k \times k$.

Considering $\mathbf{A}_n$ we have

$$S_n^n \langle \lambda_i \rangle = |\mathbf{A}_n| = 1, \quad (4)$$

$$S_1^n \langle \lambda_i \rangle = \text{tr}(\mathbf{A}_n) = \frac{n(n+1)}{2}. \quad (5)$$

A closed inspection of the structure of the determinants in Equation 3 show that

$$\begin{aligned} S_k^n \langle \lambda_i \rangle &= \sum_{i_1=1}^{n-k+1} \sum_{i_2=1}^{n-k+2-i_1} \sum_{i_3=1}^{n-k+3-i_1-i_2} \cdots \\ &\quad \cdots \sum_{i_k=1}^{n-i_1-i_2-\dots-i_{k-1}} \Delta_k(i_1, i_2, i_3, \dots, i_k) \\ &= \sum_{i_1=1}^{n-k+1} \sum_{i_2=1}^{n-k+2-i_1} \sum_{i_3=1}^{n-k+3-i_1-i_2} \cdots \\ &\quad \cdots \sum_{i_k=1}^{n-i_1-i_2-\dots-i_{k-1}} i_1 i_2 \cdots i_k. \end{aligned}$$

We can develop this formula according the index i_1 in the following way

$$\begin{aligned}
S_k^n \langle \lambda_i \rangle &= 1 \times \sum_{i_2=1}^{n-k+1} \sum_{i_3=1}^{n-k+2-i_2} \cdots \sum_{i_k=1}^{n-1-i_2-\cdots-i_{k-1}} i_2 i_3 \cdots i_k \\
&+ 2 \times \sum_{i_2=1}^{n-k} \sum_{i_3=1}^{n-k+1-i_2} \cdots \sum_{i_k=1}^{n-2-i_2-\cdots-i_{k-1}} i_2 i_3 \cdots i_k \\
&+ \cdots \cdots \\
&+ (n-k+1) \times \sum_{i_2=1}^1 \sum_{i_3=1}^{2-i_2} \cdots \sum_{i_k=1}^{k-1-i_2-\cdots-i_{k-1}} i_2 i_3 \cdots i_k.
\end{aligned}$$

This shows that we can write

$$S_k^n \langle \lambda_i \rangle = \sum_{i=1}^{n-k+1} i S_{k-1}^{n-i} \langle \lambda_i \rangle. \quad (6)$$

In the above formula we set $S_0^n \langle \lambda_i \rangle = 1$.

Using this equation we have

$$S_k^n \langle \lambda_i \rangle = S_{k-1}^{n-1} \langle \lambda_i \rangle + 2S_{k-1}^{n-2} \langle \lambda_i \rangle + 3S_{k-1}^{n-3} \langle \lambda_i \rangle + \cdots + (n-k+1)S_{k-1}^{k-1} \langle \lambda_i \rangle,$$

and

$$S_k^{n-1} \langle \lambda_i \rangle = S_{k-1}^{n-2} \langle \lambda_i \rangle + 2S_{k-1}^{n-3} \langle \lambda_i \rangle + 3S_{k-1}^{n-4} \langle \lambda_i \rangle + \cdots + (n-k)S_{k-1}^{k-1} \langle \lambda_i \rangle.$$

It follows

$$S_k^n \langle \lambda_i \rangle = S_k^{n-1} \langle \lambda_i \rangle + \sum_{i=1}^{n-k+1} S_{k-1}^{n-i} \langle \lambda_i \rangle. \quad (7)$$

This formula makes it easy to evaluate $S_k^n \langle \lambda_i \rangle$ for all n and k .

Consider the following identity relating to binomial coefficients

$$\binom{n+k}{n-k} = \binom{n+k-1}{n-k-1} + \sum_{i=1}^{n-k+1} \binom{n+k-1-i}{n-k+1-i}, \quad 0 \leq k \leq n. \quad (8)$$

If we set

$$S_k^n = \binom{n+k}{n-k},$$

we see that

$$S_k^{n-1} = \binom{n+k-1}{n-k-1},$$

and

$$S_{k-1}^{n-i} = \binom{n+k-1-i}{n-k+1-i}.$$

In this way Equation 7 follows from identity 8, and conversely. So we can conclude that

$$S_k^n \langle \lambda_i \rangle = \binom{n+k}{n-k}. \quad (9)$$

Note that

$$\binom{n+k}{n-k} = \binom{n+k}{2k}.$$

Then Identity 1.76 in [3] says

$$\sum_{k=0}^n \binom{n+k}{n-k} = F_{2n+1},$$

where F_i is the i -th Fibonacci number. It follows

$$\begin{aligned} \sum_{k=1}^n S_k^n \langle \lambda_i \rangle + 1 &= \sum_{k=1}^n \binom{n+k}{2k} + 1 \\ &= \sum_{k=0}^n \binom{n+k}{2k} \\ &= F_{2n+1}. \end{aligned} \quad (10)$$

We can get another recurrence relationship. Indeed from

$$\begin{aligned} \frac{n+k}{n-k} \binom{n+k-1}{n-k-1} &= \frac{(n+k)(n+k-1)!}{(n-k)(n-k-1)!(2k)!} \\ &= \frac{(n+k)!}{(2k)!(n-k)!} \\ &= \binom{n+k}{n-k}, \end{aligned}$$

we get

$$S_k^n \langle \lambda_i \rangle = \frac{n+k}{n-k} S_k^{n-1} \langle \lambda_i \rangle, \quad k = 1, 2, \dots, n-1 \quad (11)$$

References

- [1] M. Catalani (1996), *Spazi Lineari*, Iuculano Editore, Pavia.
- [2] F.R. Gantmacher (1977), *The Theory of Matrices*, Chelsea Publishing Company, New York.
- [3] H.W. Gould (1972), *Combinatorial Identities*, Morgantown, W. Va.
- [4] C.R. Rao and S.K. Mitra (1971), *Generalized Inverse of Matrices and its Applications*, Wiley and Sons, New York.